

XVI. *On some rare and beautiful Insects from Silhet, chiefly in the Collection of FREDERICK JOHN PARRY, Esq., F.L.S., &c. By the Rev. FREDERICK WILLIAM HOPE, M.A., F.R.S., F.L.S., &c. Continued from page 112.*

Read November 1st, 1842.

Order LEPIDOPTERA.

Family PAPILIONIDÆ.

TEINOPALPUS, *Hope*, n. g.

*Corpus robustum; capite fronte conico. Palpi capite longiores, conici, porrecti. Antennæ clavâ sensim increscenti. Pedes antici perfecti, tibiis in medio appendiculatis. Alæ anticæ apice acutè falcatæ; cellulâ discoidali clausâ; venâ medianâ ramos quatuor emittente, ramo postico venæ subcostalis ad ramum furcatum præcedentem approximato. Alæ posticæ caudatae.*

Sp. 16. TEINOPALPUS IMPERIALIS. TAB. XI. fig. 1, 2.

Alis viridi-pulverosissimis; anticis fasciâ tenui transversâ medianâ nigrâ extùs flavo-marginatâ nebulisque duâbus fuscis submarginalibus; posticis caudâ unicâ terminatis maculâ magnâ medianâ flavâ nigro-cinctâ squamulis cinnereis lunulisque marginalibus flavis viridibusque; omnibus subtùs aurantiis nigro-striatis dimidio basali viridi; posticis apicibus nigris griseo viridine variis.

Long. corp. lin. 15; expans. alar. unc. 3, lin. 10.

Habitat in Indiâ Orientali, Silhet; e Museo Dom. Parry descriptus.

Sp. 17. TEINOPALPUS PARRYÆ. TAB. XI. fig. 3, 4.

Affinis præcedenti at major, alis omnibus basi viridibus; anticis minùs falcatis nigro transversè strigatis nebulosisque; posticis bicaudatis plagâ magnâ medianâ pallidè luteâ nigro-pulverosâ strigâque undulatâ nigrâ:

extus griseo-pulverosis lunulisque marginalibus viridibus flavisque ornatis: angulo anali latè flavo.

Long. corp. lin. 17; expans. alar. unc. 4, lin. 7.

*Habitat* in Indiâ Orientali, Silhet.

The above magnificent insect was sent to me by Captain Parry for description, and I have consequently named it after Mrs. Parry: it is possible that the present may be the female of the former species. There is, I believe, only another specimen in this country, which is in the rich Lepidopterous collection of the East India House.

#### Family BOMBYCIDÆ.

##### Genus SATURNIA.

Sp. 18. SATURNIA ZULEIKA. TAB. XI. fig. 5.

Alis pallidè puniceis fusco-irroratis albidoque variis: apicibus fuscis; anticis plagâ magnâ basali rufo-brunneâ; omnibus maculâ magnâ medianâ semi-lunari sanguineo-fuscâ lunulis tribus albis griseisque lineis ternis fuscis undulatis valdè angulatis inter ocellum marginemque posticum aliisque duâbus submarginalibus.

Long. corp. lin. 14; expans. alar. unc. 4, lin. 8.

*Habitat* in Silhet.

#### Order HOMOPTERA.

##### Family FULGORIDÆ.

##### Genus APHANA.

Sp. 19. APHANA AMABILIS. TAB. XII. fig. 1.

Viridis, alis anticis cretaceis maculis numerosis ovalibus aspersis; posticis coccineis: apicibus pallidè virescentibus.

Long. corp. lin. 12; expans. alar. unc. 2, lin. 9.

*Corpus* suprà aurantium; capite viridi, cornu erecto acuto tenui. *Prothorax* viridis, maculis tribus fulvis insignitus. *Corpus* infrà viride; pedibus concoloribus; pectore lanagine albida asperso.

This remarkably beautiful insect was received lately from Silhet; the spots

which on the upper side appear to be white and green intermixed, are beneath entirely green.

Sp. 20. *APHANA AURORA.* TAB. XII. fig. 2.

Affinis *Aphanæ aurantiæ*, Hope; at minor, ferè totum corpus aurantium, alis anticis externè vix virescentibus internè aurantiis maculisque albidis minutissimis aspersis.

Long. corp. lin. 9; expans. alar. unc. 2, lin. 5.

*Habitat* in Indiâ Orientali, Silhet.

*Corpus* infrâ concolor; alis maculis cretaceis insignitis; tibiis posticis externè spinosis.

LYSTRA.

Sp. 21. *LYSTRA WESTWOODII*, Parry. TAB. XII. fig. 3.

Fusco-flava, rostro concolore inter oculos recurvo, fronte ferè trigono elongato, alis anticis dimidio basali fusco-flavis maculis fuscis magnitudine variis aspersis; posticis basi sanguineo colore fucatis lineâ atrâ marginatis punctisque tribus insignitis: reliquâ parte immaculatâ hyalinâ, corpore infrâ nigro.

Long. corp. lin. 6; expans. alar. unc. 1, lin. 11.

The above insect is described from the rich collection of Captain Parry.

Sp. 22. *LYSTRA DIMIDIATA.* TAB. XII. fig. 4.

Fusca, capite thorace pedibusque concoloribus, alis anticis dimidio basali virescente maculis minutis crebris atris asperso: apicibus subaurantio-hyalinis maculis cretaceis insignitis; posticis dimidio basali viridi: posteriori albidâ atropunctatâ: apicibus fuscis maculis virescentibus irroratis.

Long. corp. lin. 7; expans. alar. unc. 1, lin. 10.

*Habitat* in Silhet. In Museo Dom. Hope.

Sp. 23. *LYSTRA PUNICEA.* TAB. XII. fig. 5.

Punicea, capite thoraceque pallidioribus, alis anticis dimidio basali roseo colore tincto maculis crebris nigris asperso: apicibus fusco-puniceis hyalinis maculâ in medio disco pellucidâ; posticis basi puniceis maculis atris insignitis: medio albidis: apicibus fusco-hyalinis.

Long. corp. lin. 6; expans. alar. unc. 1, lin. 11.

*Habitat* in Silhet; e Museo Dom. Hope descripta.

## EURYBRACHIS.

## Sp. 24. EURYBRACHIS BASALIS. TAB. XII. fig. 6.

Fusco-flava, capite flavescente, thorace brunneo, abdomine miniato, alis anticis fuscis flavo-reticulatis; posticis ad basin antrorsum lineâ cæruleâ incurvâ insignitis posticè subroseis: reliquâ parte nigricante.

Long. corp. lin. 10; expans. alar. unc. 2, lin. 3.

*Habitat* in Silhet.

## Sp. 25. EURYBRACHIS PULVEROSA. TAB. XII. fig. 7.

Candida, capite thoraceque flavis, abdomine concolore, caudâ gossypio ornata, alis anticis dimidio basali virescente: apicibus fusco-hyalinis sparsim maculatis; posticis albidis maculâ irregulari nigrâ ad marginem anticum aliisque minutis concoloribus posticè locatis, pedibus quatuor anterioribus miniatis; posticis fuscis.

Long. corp. lin. 7; expans. alar. unc. 2, lin. 3.

*Habitat* in Silhet.

## Sp. 26. EURYBRACHIS REVERSA. TAB. XII. fig. 8.

Fusco-flava, capite thoraceque concoloribus, abdomine medio albido: posticè gossypio flavo ornato: apice candido, alis anticis fusco-flavis ad basin subsanguineo colore tinctis: maculâ albâ rotundatâ ad marginem anticum secundâ minore ferè circa medium disci aliisque minutissimis sparsim irroratis; posticis dimidio basali albido: apicibus irregulariter fusco-irroratis, pedibus quatuor anterioribus flavo-fuscis: tibiis obscurioribus; posticis femoribus pallidè testaceis: tibiis nigricantibus.

Long. corp. lin. 7; expans. alar. unc. 2, lin. 7.

*Habitat* in Indiâ Orientali, Silhet.

Sp. 27. EURYBRACHIS INSIGNIS, *Westwood*. TAB. XII. fig. 9.

Candida, capite thoraceque flavis, abdomine lateribus sanguineo colore fucatis, alis anticis flavescentibus maculis variis minutis nigris aspersis; posticis albidis maculâ irregulari nigrâ ad marginem anteriorem aliisque minutis concoloribus posticè; pedibus anterioribus fusco-flavis; posterioribus viridibus: tibiarum spinis nigricantibus.

Long. corp. lin. 7; expans. alar. unc. 2, lin. 2.

*Habitat* in regione Malabaricâ, e Museo Dom. Westwood descripta.

**CORETHRURA\***, *Hope*, n. g.

*Corpus* breve, crassum, abdominis apice longissimè floccoso. *Caput* anticè carinatum, ferè ut in genere *Derbe*; facie anticè visâ angustâ. *Promuscis* longitudine mediocris. *Clypeus* magnus, inflatus. *Antennæ* sub oculos insertæ, brevissimæ, apice longè setoso. *Ocelli* minuti, paulò ante oculos locati. *Pedes antici* tibiis depresso-instructi. *Tibiae posticæ* extùs 3-dentatæ.

Sp. 28. **CORETHRURA FUSCO-VARIA.** TAB. XII. fig. 10.

Subfusca, capite lineâ transversâ albida insignito, abdome fusco-viridi: caudâ subochraceâ gossypio ornatâ, alis anticis fusco-variis farinâ albida irroratis; posticis pallidioribus hyalinis.

Long. corp. lin. 9; expans. alar. unc. 2, lin. 4.

*Habitat* in Silhet. In Museo Dom. Hope.

---

**EXPLANATION OF THE PLATES.**

**TAB. XI.**

- Fig. 1. *Teinopalpus Imperialis*, upper side.
2. *Teinopalpus Imperialis*, under side.
3. *Teinopalpus Parryæ*, upper side.
4. *Teinopalpus Parryæ*, under side.
5. *Saturnia Zuleika*.

**TAB. XII.**

- Fig. 1. *Aphana amabilis*.
2. *Aphana Aurora*.
3. *Lystra Westwoodii*.

\* From κόρηθρον a broom, and οὐρὰ a tail.

Fig. 4. *Lystra dimidiata.*

5. *Lystra punicea.*

6. *Eurybrachis basalis.*

7. *Eurybrachis pulverosa.*

8. *Eurybrachis reversa.*

9. *Eurybrachis insignis.*

10. *Corethrura fusco-varia.*





