



First record of the family Issidae (Hemiptera, Auchenorrhyncha, Fulgoroidea) from the Hawaiian Islands

Vladimir Gnezdilov[‡], Charles R. Bartlett[§]

[‡] Zoological Institut RAS, St-Petersburg, Russia

[§] University of Delaware, Newark, United States of America

Corresponding author: Charles R. Bartlett (bartlett@udel.edu)

Academic editor: J. Adilson Pinedo-Escatel

Received: 05 Jan 2022 | Accepted: 15 Mar 2022 | Published: 22 Mar 2022

Citation: Gnezdilov V, Bartlett CR (2022) First record of the family Issidae (Hemiptera, Auchenorrhyncha, Fulgoroidea) from the Hawaiian Islands. Biodiversity Data Journal 10: e80135.

<https://doi.org/10.3897/BDJ.10.e80135>

Abstract

Background

Euroxenus vayssieresii (Bonfils, Attie & Reynaud, 2001) (Issinae, Sarimini) was described (in the genus *Borbonissus* Bonfils, Attie & Reynaud, 2001) from Réunion Island, in the Indian Ocean and, previous to this report, has not been recorded elsewhere. *Euroxenus vayssieresii* is here illustrated and re-described to improved taxonomic diagnosis.

New information

Euroxenus vayssieresii is recorded for the first time from the Island of Hawaii in the Hawaiian Archipelago. This is first record of the family Issidae from the Hawaiian Archipelago.

Keywords

Issidae, new record, Pacific Region, Sarimini, Hawaii

Introduction

The Hawaiian planthopper (Hemiptera, Auchenorrhyncha, Fulgoroidea) fauna consists of 207 endemic species (64 Cixiidae plus 143 Delphacidae) and at least 15 adventive species (10 Delphacidae, 2 Flatidae, 1 Tropiduchidae and 2 Derbidae) (Zimmerman 1948, Beardsley 1979, Beardsley 1990, Asche 1997, Asche 2000a, Asche 2000b, Nishida 2002, Hoch 2006). No additional adventive planthopper species have been reported since the publication of the Fourth Edition of Bishop Museum's Hawaiian Terrestrial Arthropod Checklist (Nishida 2002), i.e. Matsunaga et al. (2019). Beardsley (1979) reported that the rate of 'accidental immigration and establishment' was about 16 species a year between 1937 and 1961 and about 19 species a year between 1962 and 1976 (subsequent rates have not been reported). The importation of live plant material may be the main source of immigrant species (at least for phytophagous insects) (Beardsley 1979, DeNitto et al. 2015, Mound et al. 2017) and the primary sources of introductions are the mainland, the United States and Asia-Pacific (DeNitto et al. 2015). Here, we report an established species of Issidae, found at three localities on the western side of the Island of Hawaii in 2021.

The planthopper family Issidae comprises about 1,090 species in 217 genera distributed worldwide (Bourgoin 2021). Issids are absent in southern Africa (except for two species of the genus *Ikonzia* Hesse, 1925 from north Namibia and south Angola), Madagascar, the Seychelles and Tasmania (Gnezdilov 2013, Gnezdilov 2016). In Oceania, Issidae are known from New Guinea (eight genera, 17 species), the Solomon Islands (one species) and Fiji (one species) (Gnezdilov 2013, Gnezdilov et al. 2015, Gnezdilov 2020b), but are absent in Micronesia and Polynesia including New Zealand and Hawaii. In the South Indian Ocean, the Issidae are known only from the Mascarene Archipelago (two species, Bonfils et al. 2001), consisting (in part) of the large islands Mauritius, Réunion and Rodrigues.

The genus *Borbonissus* Bonfils, Attie & Reynaud, 2001 was described with two species from Réunion Island, *B. brunneifrons* Bonfils, Attie & Reynaud, 2001 (type species) and *B. vayssieresii* Bonfils, Attie & Reynaud, 2001 (Bonfils et al. 2001). Later, *Borbonissus* was synonymised under *Thabena* Stål 1866 and *Euroxenus* Gnezdilov, 2009, erected to accommodate *B. vayssieresii* (Gnezdilov 2009). Gnezdilov (2009) postulated that the issid fauna of Réunion Island had an Oriental genesis and proposed a close relationship of *Euroxenus* and the Oriental genus *Eusarima* Yang, 1994 (in Chan and Yang 1994), recently supported by molecular data (Gnezdilov et al. in press). *Thabena brunneifrons* was later recorded from Rodrigues, Taiwan, Dongsha (Pratas) Islands and Singapore (Chan et al. 2013, Gnezdilov 2014b, Gnezdilov 2015).

Here, we record *E. vayssieresii* from the Island of Hawaii in the Hawaiian Archipelago. This species is re-described to improve on diagnostic features reported by Gnezdilov (2009) and Gnezdilov (2020).

Materials and methods

Morphological terminology follows Anufriev and Emeljanov (1988) and Gnezdilov et al. (2014). Photographs were taken using a Canon EOS 5D Mark IV camera with the lens Canon-MP-E-65mm f/2,8 1-5x Macro and the flash Canon Macro Twin-Lite MT-26EX-RT. Images were produced using Helicon Focus v. 7.6.4 and Adobe Photoshop CC 2019 software. The genital segments of male specimens examined were macerated in 10% potassium hydroxide (KOH) and figured in glycerine jelly (Brunel Micro Ltd, UK) using a Leica MZ9.5 stereomicroscope with a camera lucida. The map was prepared using Google Earth Pro (version 7.3) with map data sources as attributed in image.

The specimens examined are deposited in University of Delaware, Department of Entomology and Wildlife Ecology, Newark, Delaware, USA (**UDCC**); Smithsonian Institution, National Museum of Natural History, Washington, DC (**USNM**); Bernice P Bishop Museum of Natural History, Honolulu, Hawaii (**BPBM**); and the Zoological Institute of the Russian Academy of Sciences, Saint Petersburg, Russian Federation (**ZIN**).

Taxon treatment

Euroxenus vayssieresii (Bonfils, Attié & Reynaud, 2001)

- GBIF <https://www.gbif.org/species/2050778>

Nomenclature

= *Borbonissus vayssieresii* Bonfils, Attié & Reynaud, 2001: 220 (original description).

= *Euroxenus vayssieresii* (Bonfils, Attié & Reynaud, 2001), combination by Gnezdilov 2009: 84.

Type locality. Réunion, la Possession, Chemin des Anglais.

Materials

- scientificName: *Euroxenus vayssieresii*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 13-08-2021; individualCount: 3; sex: Male; lifeStage: adult; catalogNumber: UDCC_TCN 00097423; recordedBy: W.D. Pereira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen
- scientificName: *Euroxenus vayssieresii*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 13-08-2021; individualCount: 3; sex: Female; lifeStage: adult; catalogNumber: UDCC_TCN 00097423; recordedBy: W.D. Pereira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen

- c. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 13-08-2021; individualCount: 1; lifeStage: nymph; catalogNumber: UDCC_TCN 00097423; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen
- d. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 16-07-2021; individualCount: 5; sex: Male; lifeStage: adult; catalogNumber: UDCC_TCN 00101252; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC, ZIN; basisOfRecord: PreservedSpecimen
- e. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 03-07-2021; individualCount: 8; sex: Male; lifeStage: adult; catalogNumber: UDCC_TCN 00101401; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC, USNM, BPBM; basisOfRecord: PreservedSpecimen
- f. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 03-07-2021; individualCount: 5; sex: Female; lifeStage: adult; catalogNumber: UDCC_TCN 00101405; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC, ZIN; basisOfRecord: PreservedSpecimen
- g. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 16-07-2021; individualCount: 4; sex: Female; lifeStage: adult; catalogNumber: UDCC_TCN 00101414; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC, ZIN; basisOfRecord: PreservedSpecimen
- h. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Keauhou Senic Lookout; verbatimElevation: 91 m; decimalLatitude: 19.57611; decimalLongitude: -155.95917; georeferenceProtocol: label; samplingProtocol: Sticky Trap; eventDate: 10/14-05-2021; individualCount: 1; sex: Male; lifeStage: adult; catalogNumber: UDCC_TCN 00101417; recordedBy: W.D. Perreira, N.G. Miller, and D.A. Yee; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen
- i. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: HonoKohau Harbor; verbatimElevation: 3 m; decimalLatitude: 19.67111; decimalLongitude: -156.02056; georeferenceProtocol: label; samplingProtocol: Sticky Trap; eventDate: 02/16-07-2021; individualCount: 1; sex: Female; lifeStage: adult; catalogNumber: UDCC_TCN 00101418; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen

- j. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 03-07-2021; individualCount: 1; sex: Male; lifeStage: adult; catalogNumber: UDCC_TCN 00101436; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen
- k. scientificName: *Euroxenus vayssieres*; country: USA; stateProvince: Hawaii; county: Hawaii; locality: Spencer Beach Park; verbatimElevation: 12 m; decimalLatitude: 20.02278; decimalLongitude: -155.82111; georeferenceProtocol: label; samplingProtocol: Sweep Netting; eventDate: 03-07-2021; individualCount: 1; sex: Female; lifeStage: adult; catalogNumber: UDCC_TCN 00101437; recordedBy: W.D. Perreira; identifiedBy: V.M. Gnezdilov; dateIdentified: 2021; ownerInstitutionCode: UDCC; basisOfRecord: PreservedSpecimen

Description

Colouration. General colouration light brown-yellowish to dark brown on anteclypeus and forewing clavus (Figs 1, 2, 3, 4). Pedicel, apices of fore and middle tibiae and apices of spines black. Forewings with black marginal cells along the costal and lateral margins (Figs 1, 2) and sometimes (Fig. 4) with dark brown band from the costal margin to inner claval margin in basal half on females. Hind femora and outer surfaces of hind tibiae dark brown. Hind margins of gonostyli dark brown to black on dorsolateral angles. Ungues dark brown to black. Gonoplacs dark brown.



Figure 1. [doi](#)

Euroxenus vayssieres, Hawaii Island, male dorsal view.




Figure 2. [doi](#)

Euroxenus vayssieresi, Hawaii Island, male left lateral view.



Figure 3. [doi](#)

Euroxenus vayssieresi, Hawaii Island, male frontal view.

Structure. Body length, males and females – 4.0 mm. Metope (~ frons, Fig. 3) broad, widest below the eyes, with distinct median carina running from upper margin through postclypeus; sublateral carinae distinct, joined with median carina at metopial upper margin (carinae somewhat projecting along upper margin), almost reaching metopoclypeal suture. Postclypeus broadly rhomboid. Rostrum reaching hind coxae, 2nd and 3rd segments are almost equal in length; 3rd one narrowing apically. Antennal scape very short, pedicel globular, just longer than wide. Coryphe (~ vertex, Figs 1, 4) as concave hexagon, slightly wider than long medially, anterior margin produced, obtusely angled, posterior margin concave. Metope and coryphe (in lateral view, Fig. 2) joined at obtuse angle (dorsal portion of sublateral carinae somewhat projecting). Ocelli vestigial. Eyes large, each eye nearly as wide as coryphe.



Figure 4. [doi](#)

Euroxenus vayssieresii, Hawaii Island, female dorsal view.

Pronotum slightly shorter than coryphe (Figs 1, 4), with smooth median carina, anterior margin strongly convex, posterior margin nearly straight, with weak median concavity. Paradiscal fields very narrow behind eyes, paranotal lobes flat, apically rounded. Mesonotum length 1.5 times pronotum at mid-line, with median and lateral carinae. Tegulae large. Forewing slightly narrowing apically (Bonfils et al. 2001, fig. 22) (Figs 1, 2, 4), without hypocostal plate and with wide precostal area; clavus long, 4/5 of wing length, broadly triangular. Forewing vein sequence: R 2, furcating near basal cell; M 2, furcating after wing middle; CuA 2, furcating after wing middle; basal cell narrowly oval.

Hind wing with coupling lobe and deep cubital and vannal clefts (Fig. 5); anal lobe less wide than remigial and remigio-vannal lobes. Hind wing vein sequence: R 2, furcating apically after coupling lobe; r-m 1; M 1; m-cua 1; CuA 2; CuP 1; Pcu 1; A₁ 2; A₂ 1. CuA₂ and CuP fused and flattened apically and Pcu and A_{1.1} fused medially. Front and middle femora slightly flattened. Hind tibia with two large lateral spines past mid-length and six apical spines. First and second metataromeres of subequal length, but basitarsus wider; basitarsus with two latero-apical spines and 13 intermediate spines arranged in arc; ventral surface with long setae; second metatarsomere with pair of latero-apical spines and median lobe.




Figure 5. [doi](#)

Euroxenus vayssièresi hindwing venation. Abbreviations: R = radius, M = media, CuA = anterior cubitus, CuP = posterior cubitus, Pcu = postcubitus, A1.1 = anterior branch of first anal vein; A1.2 = posterior branch of first anal vein; A2 = second anal vein.

Male terminalia: Pygofer (in lateral view, Fig. 6A) elongate vertically, with convex hind margins. Hind margin of sternite VII straight (ventral view). Phallobase (Fig. 6B and C) narrow and strongly curved (in lateral view), with a narrow process dorsally (Fig. 6B *dp*); each dorsolateral phallobase lobe with three-branched large process – two branches directed dorsally and the longest branch directed basally, with serrulate carina. Each dorsolateral phallobase lobe (in lateral view) with a lateral slit and folds near middle and with narrow semicircular subapical lobe. Ventral phallobase lobe long and wide, with an apical notch. Aedeagus with a pair of long (0.3× length of phallobase, Fig. 6B and C), narrow, apically pointed ventral hooks, arising subapically and directed basally. Apical aedeagal processes long and wide apically (Fig. 6B, shaded). Gonostylus (Fig. 6E) with deeply concave hind margin. Capitulum wide, not narrowing apically (in dorsal view, Fig. 6F), with small lateral tooth (in lateral view). Connective (lateral view, Fig. 6B) with small cup and long handle, in caudal view (Fig. 6G), with ventral margin bilobed (these articulating with the gonostyli). Anal tube elongate (Fig. 6A), in dorsal view (Fig.

6D) nearly 3 times as long as wide medially, slightly narrowing apically, with weak apical concavity. Anal column 0.3x as long as anal tube, narrow.




Figure 6. [doi](#)

Euroxenus vayssièresi, male terminalia (A–B, D–G – Hawaii; C – Réunion Island, La Possession) **A** pygofer and anal tube, lateral view; **B** penis and connective (apical aedeagal processes shaded), lateral view; **C** penis (apical aedeagal processes not visible), lateral view; **D** anal tube, dorsal view; **E** gonostylus, lateral view; **F** gonostylus, dorsal view; **G** connective handle, dorso-caudal view. Abbreviation: dp – dorsal process of the phallobase.

Female terminalia: Hind margin of sternite VII protruding medially (ventral view). Gonoplasts short. Anal tube elongate. Anal column short, 0.25 of anal tube length, narrow.

Notes: *Gnezdilov (2020)* (in fig. 2) had not observed a small basal process of three-branched processes of dorsolateral phallobase lobes (Fig. 6B and C) on his drawings of male genitalia of *E. vayssièresi* from Réunion Island (Saint-Paul); otherwise no significant differences in external morphological features were found between the specimens from Réunion (La Possession and Saint-Paul) and those of the Hawaii Islands.

Euroxenus Gnezdilov, 2009

Genus diagnosis: Metope (~ frons) broad (Fig. 3), widest below the eyes, with distinct median carina running from upper margin through postclypeus and distinct sublateral

carinae joined with median carina at upper metopial margin and almost reaching metopoclypeal suture. Rostrum reaching hind coxae, 2nd and 3rd segments are almost equal in length; 3rd narrowing apically. Coryphe (~ vertex) (Fig. 4) slightly wider than long medially, anterior margin produced, obtusely angled. Metope and coryphe joined at obtuse angle (in lateral view, Fig. 2). Paradiscal fields of pronotum very narrow behind eyes. Forewings just covering abdominal apex, slightly narrowing apically, without hypocostal plate, with wide precostal area; clavus long. Forewing vein sequence: R 2, furcating near basal cell; M 2, CuA 2, both furcating after wing middle. Hind wings (Fig. 5) with deep cubital and vannal clefts; anal lobe less wide than remigial and remigio-vannal lobes. Hind wing vein sequence: R 2; r-m 1; M 1; m-cua 1; CuA 2; CuP 1; Pcu 1; A₁ 2; A₂ 1. CuA and CuP fused and flattened apically and Pcu and A₁ fused medially. Hind tibia with two large lateral spines. First metataromere with two latero-apical spines and 13 intermediate spines arranged in arc. Phallobase narrow, curved (in lateral view, Fig. 6B and C), each dorsolateral phallobase lobe with narrow semicircular subapical lobe, large three-branched process below it and with a lateral slit and folds near middle. Ventral phallobase lobe long and wide. Aedeagus with a pair of long and narrow, apically pointed ventral hooks, arising subapically and directed basally. Apical aedeagal processes long and wide apically. Gonostylus (in lateral view, Fig. 6E) with deeply concave hind margin. Hind margin of female sternite VII protruding medially (ventral view); gonoplags short; anal tube elongate.

Differential genus diagnosis: *Euroxenus* is closely related to *Eusarima* Yang, 1994 and *Duplexissus* Wang, Zhang & Bourgoïn, 2019, based on having completely developed median and sublateral carinae of metope and the presence of a long and narrow process on each dorso-lateral lobe of the phallobase. However, *Euroxenus* differs from both genera by the shorter forewings with bifurcate median veins (tri- or tetrafurcate in *Eusarima* and *Duplexissus*) and by trifurcate processes of the phallobase (simple in *Eusarima* and bifurcate in *Duplexissus*) (Chan and Yang 1994, figs. 45–72; Wang et al. 2019, figs. 12–22).

Discussion

The specimens of *Euroxenus vayssièresi* were all collected in 2021 in very dry areas, on the west side of the Island of Hawaii at three localities (Fig. 7, from north to south): Keauhou, Kailua-Kona and Spencer Beach Park. The three localities are all near the coast (Keauhou Lookout is about ca. 300 feet (ca. 100 m) elevation), with the straight-line distance between Keauhou and Spencer Beach about 30 miles (48 km). The dominant vegetation at the sites was exotic species with a Kiawe (*Prosopis pallida*) and Koa Haole (*Leucaena leucocephala*) canopy. The specimens were mainly collected by general sweeping of grasses and weeds of the understorey, with no clear plant associations made.

In Réunion, adults and nymphs of *E. vayssièresi* were recorded from *Cestrum* sp. (stated as 'noctuiiflorum', but apparently meaning *nocturnum*, Bourgoïn 2021) and *Capsicum* sp. (Solanaceae), *Bridelia micrantha* (Phyllanthaceae) and *Coccoloba uvifera* (Polygonaceae) suggesting that this species is polyphagous (Bonfils et al. 2001, Attié et al. 2008). Bonfils et

al. (2001) noted that their specimens were collected in semi-dry secondary vegetation on exotic and native plants located on the northwest and west of the "leeward" region of Réunion, which is similar to the circumstances where this species was found in Hawaii.

The genus *Euroxenus* (Issinae, Sarimini sensu Wang et al. 2016 and Gnezdilov et al. 2020) was monotypic as described. Gnezdilov (2020) placed the monotypic genus *Duplexissus* Wang, Zhang & Bourgoïn, 2019 (Wang et al. 2019) in synonymy under *Euroxenus*, but this placement was disputed by Wang et al. (2020) on both morphological and molecular grounds.




Figure 7. [doi](#)

Collecting localities for *Euroxenus vayssieresii* on the Island of Hawaii (black circles). Map created using Google Earth, map data sources provided in the lower left of the image.

Issids are not typically good dispersers (many are sub-brachypterous and flightless) and transport of *Euroxenus vayssieresii* to Hawaii is most likely by human agency and in association with live plants. This was previously hypothesised as the mode of transport for other issid species (Gnezdilov and O'Brien 2006, Gnezdilov 2009).

Acknowledgements

We thank William D. Perreira, CEO (Chief Entomological Officer) of the Dana Anne Yee Foundation for collecting and bringing this species to our attention. We thank The Atherton Family Foundation and The Dana Anne Yee Foundation for funding of collecting and the County of Hawaii, Department of Parks and Recreation for permission to collect at Spencer Beach Park. We are grateful to Dr. Vladimir V. Neimorovets (Saint Petersburg, Russia) for taking photos of the species. The study of VMG was performed, based on the ZIN

collection within the State Program no. 1021051302540-6. Research by CRB was partially supported by USDA Hatch (DEL00774).

References

- Anufriev GA, Emeljanov AF (1988) Suborder Cicadinea (Auchenorrhyncha). In: Ler PA (Ed.) Keys to the insects of the Far East of the USSR. Vol. 2. Nauka, Leningrad, Russia, 12–495 pp. [In Russian].
- Asche M (1997) A review of the systematics of Hawaiian planthoppers (Hemiptera: Fulgoroidea). *Pacific Science* 51: 366-376. URL: <http://hdl.handle.net/10125/3194>
- Asche M (2000a) New state records of immigrant planthoppers in Hawaii (Homoptera: Fulgoroidea). *Proceedings of the Hawaiian Entomological Society* 43: 205-207. URL: <http://hdl.handle.net/10125/8409>
- Asche M (2000b) *Emoloana*, a new genus for the endemic grass-feeding Hawaiian Delphacidae (Homoptera Fulgoroidea). *Proceedings of the Hawaiian Entomological Society* 34: 71-94. URL: <http://hdl.handle.net/10125/8418>
- Attié M, Bourgoïn T, Veslot J, Soulier-Perkins A (2008) Patterns of trophic relationships between planthoppers (Hemiptera: Fulgoromorpha) and their host plants on the Mascarene Islands. *Journal of Natural History* 42: 1591-1638. <https://doi.org/10.1080/00222930802106963>
- Beardsley JW (1979) New immigrant insects in Hawaii: 1962 through 1976. *Proceedings of the Hawaiian Entomological Society* 23: 35-44. URL: <http://hdl.handle.net/10125/11090>
- Beardsley JW (1990) Notes on immigrant delphacid planthoppers in Hawaii (Homoptera: Fulgoroidea). *Proceedings of the Hawaiian Entomological Society* 30: 121-129. URL: <http://hdl.handle.net/10125/11270>
- Bonfils J, Attié M, Reynaud B (2001) Un nouveau genre d'Issidae de l'île de la Reunion: *Borbonissus* n. gen. (Hemiptera, Fulgoromorpha). *Bulletin de la Société Entomologique de France* 106: 217-224. [In French]. <https://doi.org/10.3406/bsef.2001.16756>
- Bourgoïn T (2021) FLOW (Fulgoromorpha Lists on The Web): a world knowledge base dedicated to Fulgoromorpha. <http://hemiptera-databases.org/flow/>. Accessed on: 2021-11-14.
- Chan M, Yang C (1994) Issidae of Taiwan: (Homoptera: Fulgoroidea). National Chung Hsing University, Department of Entomology, Taichung, Taiwan, 168 pp. [ISBN 9570908750]
- Chan M, Yeh H, Gnezdilov VM (2013) *Thabena brunneifrons* (Hemiptera: Issidae), new alien species in Taiwan, with notes on its biology and nymphal morphology. *Formosan Entomologist* 33: 149-159. URL: <http://delphacid.s3.amazonaws.com/9973.pdf>
- Chan ML, Yang CT (1994) Issidae of Taiwan. Fulgoroidea). Chen Chung Book, Taichung, 16 pp.
- DeNitto GA, Cannon P, Eglitis A, Glaeser JA, Maffei H, Smith S (2015) Risk and pathway assessment for the introduction of exotic insects and pathogens that could affect Hawai'i's native forests. U.S. Department of Agriculture, Forest Service, Pacific South-west Research Station. Gen. Tech. Rep. PSW-GTR-250. 171 pp. <https://doi.org/10.2737/PSW-GTR-250>

- Gnezdilov VM, O'Brien LB (2006) *Hysteropterum severini* Caldwell & DeLong, 1948, a synonym of *Agalmatium bilobum* (Fieber, 1877) (Hemiptera: Fulgoroidea: Issidae). Pan-Pacific Entomologist 82: 50-53. URL: <http://delphacid.s3.amazonaws.com/1800.pdf>
- Gnezdilov VM (2009) Revisionary notes on some tropical Issidae and Nogodinidae (Hemiptera: Fulgoroidea). Acta Entomologica Musei Nationalis Pragae 49: 75-92. URL: <http://delphacid.s3.amazonaws.com/5210.pdf>
- Gnezdilov VM (2013) Modern classification and the distribution of the family Issidae Spinola (Homoptera, Auchenorrhyncha, Fulgoroidea). Entomologicheskoe obozrenie 92 (4): 724-738. [In Russian]. <https://doi.org/10.1134/S0013873814050054>
- Gnezdilov VM (2014) Review of Indochinese Issini Spinola, 1839 (Hemiptera, Fulgoroidea, Issidae), with description of a new genus from Laos. Zoosystema 36: 761-770. <https://doi.org/10.5252/z2014n4a4>
- Gnezdilov VM, Holzinger WE, Wilson MR (2014) The Western Palaearctic Issidae (Hemiptera, Fulgoroidea): an illustrated checklist and key to genera and subgenera. Proceedings of the Zoological Institute RAS 318 (Supplement 1): 1-124. URL: https://www.zin.ru/journals/trudyzin/doc/vol_318_s1/TZ_318_1_Supplement_Gnezdilov.pdf
- Gnezdilov VM (2015) First coloured species of the genus *Thabena* Stål (Hemiptera, Fulgoroidea, Issidae) from Vietnam with general notes on the genus. Acta Zoologica Academiae Scientiarum Hungaricae 61: 329-339. <https://doi.org/10.17109/azh.61.4.329.2015>
- Gnezdilov VM, Cesne ML, Soulier-Perkins A, Bourgoïn T (2015) New Guinean Issidae: description of new taxa in a poorly known island fauna (Hemiptera, Fulgoroidea). Zootaxa 3904 (1): 82-94. <https://doi.org/10.11646/zootaxa.3904.1.4>
- Gnezdilov VM (2016) A review of the genus *Ikonza* Hesse with notes on evolution of the family Issidae (Hemiptera: Auchenorrhyncha: Fulgoroidea). Entomological Review 96: 225-234. <https://doi.org/10.1134/S0013873816020093>
- Gnezdilov VM (2020a) (dated 2019) On the synonymy and distribution of the genera *Euroxenus* Gnezdilov, 2009 and *Nikomiklukha* Gnezdilov, 2010 (Hemiptera, Auchenorrhyncha, Fulgoroidea: Issidae). Entomological Review 99: 1299-1303. <https://doi.org/10.1134/S0013873819090070>
- Gnezdilov VM (2020b) New tribe, new genera, and new species of the family Issidae (Hemiptera: Fulgoroidea) from Southeastern Asia and New Guinea, with notes on morphology and evolution of the family. Proceedings of the Zoological Institute RAS 324 (3): 306-345. <https://doi.org/10.31610/trudyzin/2020.324.3.306>
- Gnezdilov VM, Konstantinov FV, Bodrov SY (2020) New insights into the molecular phylogeny and taxonomy of the family Issidae (Hemiptera: Auchenorrhyncha: Fulgoroidea). Proceedings of the Zoological Institute RAS 324: 146-161. <https://doi.org/10.31610/trudyzin/2020.324.1.146>
- Gnezdilov VM, Konstantinov FV, Namyatova A (in press) From modern to classic: classification of the planthopper family Issidae (Hemiptera, Auchenorrhyncha, Fulgoroidea) derived from the total-evidence phylogeny. Systematic Entomology.
- Hoch H (2006) Systematics and evolution of *Iolania* (Hemiptera: Fulgoromorpha: Cixiidae) from Hawai'i. Systematic Entomology 31: 302-320. <https://doi.org/10.1111/j.1365-3113.2005.00312.x>
- Matsunaga JN, Howarth FG, Kumashiro BR (2019) New state records and additions to the alien arthropod fauna in the Hawaiian Islands. Proceedings of the Hawaiian Entomological Society 51: 1-71. URL: <http://hdl.handle.net/10125/61800>

- Mound L, Matsunaga J, Bushe B, Hoddle M, Wells A (2017) Adventive Thysanoptera species in the Hawaiian Islands: new records and putative host associations. *Proceedings of the Hawaiian Entomological Society* 49: 17-28. URL: <http://hdl.handle.net/10125/48463>
- Nishida GM (2002) Hawaiian terrestrial arthropod checklist. *Bishop Museum Technical Reports* 22: 1-313.
- Wang M, Zhang Y, Bourgoin T (2019) On the tribe Sarimini with two new genera from south of China (Hemiptera, Fulgoromorpha, Issidae). *Zootaxa* 4706: 375-383. <https://doi.org/10.11646/zootaxa.4706.2.10>
- Wang M, Zhang Y, Bourgoin T (2020) Notes on genera of Sarimini with description of two new species in the genera *Microsarimodes* and *Tetrichina* (Hemiptera: Fulgoromorpha: Issidae). *Zootaxa* 4858: 589-600. <https://doi.org/10.1016/j.ympev.2016.08.012>
- Wang ML, Zhang YL, Bourgoin T (2016) Planthopper family Issidae (Insecta: Hemiptera: Fulgoromorpha): Linking molecular phylogeny with classification. *Molecular Phylogenetics and Evolution* 105: 224-234. <https://doi.org/10.1016/j.ympev.2016.08.012>
- Zimmerman EC (1948) *Insects of Hawaii*. Volume 4. Homoptera: Auchenorrhyncha. University of Hawaii Press, Honolulu, vii+268 pp.