

Taxonomic study of the genus *Kuvera* Distant, 1906 (Hemiptera, Fulgoromorpha, Cixiidae) with descriptions of two new species from China

Yan Zhi¹, Lin Yang^{2,3}, Xiang-Sheng Chen^{2,3}

1 Key Laboratory of Medical Insects, Guizhou Medical University, Guiyang, Guizhou, 550025, China
2 Institute of Entomology, Guizhou University, Guiyang, Guizhou, 550025, China **3** The Provincial Special Key Laboratory for Development and Utilization of Insect Resources of Guizhou, Guizhou University, Guiyang, Guizhou, 550025, China

Corresponding author: Xiang-Sheng Chen (chenxs3218@163.com)

Academic editor: Pavel Stoev | Received 23 March 2022 | Accepted 13 December 2022 | Published 18 January 2023

<https://zoobank.org/917B8619-82C6-444D-A394-8D0E951DE940>

Citation: Zhi Y, Yang L, Chen X-S (2023) Taxonomic study of the genus *Kuvera* Distant, 1906 (Hemiptera, Fulgoromorpha, Cixiidae) with descriptions of two new species from China. ZooKeys 1141: 41–63. <https://doi.org/10.3897/zookeys.1141.84211>

Abstract

Two new species of genus *Kuvera* Distant, 1906, *Kuvera campyloptropa* Zhi & Chen, **sp. nov.** and *K. elongata* Zhi & Chen, **sp. nov.**, and a new Chinese record, *K. basarukini* Emeljanov, 1998, are described and illustrated from China. The females of two other species of *Kuvera*, *K. laticeps* (Metcalf, 1936) and *K. ussuriensis* (Vilbaste, 1968), are described for the first time. An updated identification key to Chinese species of *Kuvera* is given.

Keywords

Auchenorrhyncha, Eastern Palearctic region, Oriental region, planthopper, Sino-Japanese region, taxonomy

Introduction

The planthopper genus *Kuvera* was established by Distant (1906), with *K. semihyalina* Distant, 1906 as the type species by original designation. This genus belongs to the tribe Semonini of subfamily Cixiinae (Hemiptera: Cixiidae). Diagnostic features of Semonini include that the postclypeus is swollen, its clypeofrontal suture is convex, and the median carina of frons is incomplete or obscure (Holzinger et al. 2002; Emeljanov

2002). Previously 25 species in this genus have been recorded successively from Eastern Palearctic, Sino-Japanese and Oriental regions (e.g., Distant 1906; Matsumura 1914; Muir 1922; Dlabola 1957; Vilbaste 1968; Anufriev 1987; Emeljanov 1998; Tsaour et al. 1991; Rahman et al. 2017; Luo et al. 2019; Bourgoïn 2022). The latest taxonomic works on *Kuvera* by Luo et al. (2019) included a description of two new species from China, a checklist of species and an identification key to 13 Chinese species, which were useful additions to the knowledge of the Chinese fauna.

The present study of Chinese specimens has found two new species and a new Chinese record. Females of two species, *K. laticeps* (Metcalf, 1936) and *K. ussuriensis* (Vilbaste, 1968), are also described for the first time.

Materials and methods

Morphological terminology follows Bourgoïn (1987) for male genitalia, Bourgoïn et al. (2015) for wing venation and Bourgoïn (1993) for female genitalia. Body length was measured from apex of vertex to tip of forewing; vertex length represents the median length of the vertex (from the apical transverse carina to the tip of basal emargination). Fuchsin staining was used to highlight female genitalia structures studied. External morphology and drawings were done with the aid of a Leica MZ 12.5 stereomicroscope. Photographs were taken with KEYENCE VHX-6000 system. Illustrations were scanned with a CanoScan LiDE 200 and imported into Adobe Photoshop 7.0 for labeling and plate composition. The dissected male and female genitalia are preserved in glycerin in small plastic tubes pinned together with the specimens. Zoogeographic regionalization scheme follows Holt et al. (2013). The distribution map was prepared with SimpleMapp (Shorthouse 2010).

The type specimens are deposited in the Institute of Entomology, Guizhou University, Guiyang, Guizhou Province, China (**GUGC**).

Taxonomy

Genus *Kuvera* Distant, 1906

Kuvera Distant, 1906: 261; Tsaour et al. 1991: 50; Anufriev and Emeljanov 1988: 443; Emeljanov 1998: 133; Luo et al. 2019: 46.

Latoliarus Dlabola, 1957: 271; synonymized by Emeljanov 1998: 133.

Type species. *Kuvera semihyalina* Distant, 1906, original designation.

Diagnosis. For the diagnosis of *Kuvera* see Luo et al. (2019: 137).

Remarks. This genus is easily separated from other members in Semonini by the following character combinations: head including eyes narrower than pronotum; vertex short, wider than long, anterior margin of vertex obscure, with only residual

traces; vertex narrowest at subapical carina, widening towards anterior and posterior margins; anterior and posterior margins wide and parabolic, almost parallel; frons prominent, median carina only distinct on basal portion, not reaching the anterior margin of vertex; clypeus swollen, postclypeus with prominent median carina, anteclypeus carina sharp or arcuate; rostrum just reaching hind coxae; forewings with ScP+R usually forked distad of CuA, RP 3-branched, MP with 4 or 5 terminals, CuA 2 or 3-branched, and with 10–11 apical cells; metatibiotarsal formula: 6/7/(7–8); pygofer with a triangular medioventral process; aedeagus with 2 spinose processes arising near base of endosoma, and endosoma with 1–2 spinose processes; periandrium almost flat and widened at base; ovipositor elongate, orthopteroid and apically curved upwards; abdominal 9th tergite with a distinct and elliptic wax plate.

Distribution. China, Korea, Japan, (Eastern) Russia, India, Myanmar, Afghanistan.

Key to the known species (males) of *Kuvera* from China (revised from Anufriev 1987 and Luo et al. 2019)

- 1 Forewing crossed before middle by a curved, slightly broken macular fuscous fascia (Distant 1906: fig. 117)..... ***K. semihyalina* Distant, 1906**
- Forewing without fascia before middle..... **2**
- 2 Pronotum white..... ***K. longipennis* Matsumura, 1914**
- Pronotum yellow to dark brown **3**
- 3 Spinose process of endosoma long, beyond the apex of the endosoma (Fig. 5H–J) ***K. elongata* sp. nov.**
- Spinose process of endosoma not beyond the apex of the endosoma **4**
- 4 One or both of the spinous processes on lateral sides of the periandrium curved to the opposite side over its dorsal surface..... **5**
- Neither of the spinous processes on lateral sides of the periandrium curved to the opposite side over its dorsal surface **13**
- 5 Both spinose processes on lateral sides of the periandrium curved to the opposite side..... **6**
- Only one of the two spinose processes on lateral sides of the periandrium curved to the opposite side..... **7**
- 6 Spinose process on right side of periandrium strongly curved, apex directed left-ventrocaudally (Anufriev 1987: figs 69, 70) ***K. toroensis* Matsumura, 1914**
- Spinose process on right side of periandrium slightly curved, apex directed left-dorsocephally (Tsaur et al. 1991: fig. 28) ***K. transversa* Tsaur & Hsu, 1991**
- 7 Spinose process on right side of periandrium curved to left side (Fig. 6H–K) ***K. laticeps* (Metcalf, 1936)**
- Spinose process on left side of periandrium curved to right side..... **8**
- 8 Left spinose process of periandrium S-shaped **9**
- Left spinose process of periandrium not S-shaped **11**

- 9 Anal segment with apical lobes symmetrical (Tsaur et al. 1991: fig. 30)
 ***K. hama* Tsaur & Hsu, 1991**
- Anal segment with apical lobes asymmetrical **10**
- 10 Apex of left process reaching base of periandrium; endosoma process reaching
 apex of sclerotized portion of endosoma (Luo et al. 2019: figs 10, 22)
 ***K. huoditangensis* Luo, Liu & Feng, 2019**
- Apex of left process not reaching base of periandrium; endosoma process
 reaching middle of membranous portion of endosoma (Anufriev 1987:
 figs 20–22)..... ***K. vilbastei* Anufriev, 1987**
- 11 Endosoma process long, longer than two-thirds of the left spinose process of
 periandrium (Luo et al. 2019: figs 43, 44)
 ***K. longwangshanensis* Luo, Liu & Feng, 2019**
- Endosoma process short, shorter than half of the left spinose process of peri-
 andrium..... **12**
- 12 Anal tube more or less parallel-sided in dorsal view; apex of left spinose pro-
 cess of periandrium exceeding right lateral margin of periandrium (Fig. 3 F,
 J, K)..... ***K. basarukini* Emeljanov, 1998**
- Anal tube widened in the middle in dorsal view; apex of left spinose process
 of periandrium not reaching right lateral margin of periandrium (Anufriev
 1987: figs 42–45)..... ***K. flaviceps* (Matsumura, 1900)**
- 13 Right process of periandrium originated ventral surface..... **14**
- Right process of periandrium originated right side..... **15**
- 14 Anal segment with apical lobes symmetrical (Anufriev 1987: fig. 67)
 ***K. ussuriensis* (Vilbaste, 1968)**
- Anal segment with apical lobes asymmetrical (Rahman et al. 2017;
 Fig. 6F)..... ***K. yecheonensis* Rahman, Kwon & Suh, 2017**
- 15 The two spinous processes of periandrium nearly equal in length..... **16**
- The two spinous processes of periandrium not of equal length..... **18**
- 16 In lateral view, two spinose processes of periandrium arched (Tsaur et al.
 1991: fig. 25)..... ***K. taiwana* Tsaur & Hsu, 1991**
- In lateral view, two spinose processes of periandrium almost straight **17**
- 17 In dorsal view, two spinose processes of periandrium curved inwards (Anu-
 friev 1987: fig. 50) ***K. kurilensis* Anufriev, 1987**
- In dorsal view, two spinose processes of periandrium almost straight (Anu-
 friev 1987: fig. 57) ***K. tappanella* Matsumura, 1914**
- 18 Right process of periandrium longer than left one (Fig. 4H–K)
 ***K. campylotropa* sp. nov.**
- Left process of periandrium longer than right one..... **19**
- 19 Ventral base of periandrium triangular (Tsaur et al. 1991: fig. 29)
 ***K. communis* Tsaur & Hsu, 1991**
- Ventral base of periandrium roundly concaved (Tsaur et al. 1991: fig. 27)
 ***K. similis* Tsaur & Hsu, 1991**

***Kuvera basarukini* Emeljanov, 1998**

Figs 1A, B, 3

Kuvera basarukini Emeljanov, 1998: 133.

Material examined. CHINA: 1♂, Guizhou Province, Duyun City, Doupengshan (26°21'N, 107°23'E), 19 August 2017, leg. Liang-Jing Yang; 1♂, Guizhou Province, Rongjiang county, Xiaodanjiang (660–800 m) (26°20'N, 108°21'E), 13–14 September 2005, leg. Bin Zhang, Zi-Zhong Li.

Redescription. Body length: male: 5.5–5.9 mm ($N = 2$).

Coloration. General color blackish brown (Figs 1A, B, 3A, B). Eyes brown, ocelli yellowish brown. Vertex brown, pronotum dark brown and mesonotum blackish brown. Frons generally yellowish brown, blackish brown above frontoclypeal suture. Clypeus blackish brown. Rostrum generally brown except darker tip. Forewing semi-translucent, with a small irregular blackish brown spot at branch of Y-vein, stigma blackish brown. Hind tibiae yellowish brown and abdominal sternites blackish brown.

Head and thorax. Vertex (Fig. 3A) broad, 2.2 times wider than long; anterior margin arched convex, posterior margin arched concave; median carina reaching transverse carinae, indistinct. Frons (Fig. 3B) 1.2 times as wide as long, median carina indistinct, extending from slightly above level of lateral ocelli to median ocellus. Clypeus with median carina distinct and elevated throughout. Pronotum (Fig. 3A) 2.2 times longer than vertex, posterior margin nearly at right angle. Mesonotum 1.6 times longer than pronotum and vertex combined. Forewing (Fig. 3C) 3.0 times longer than wide, with 10 apical and 6 subapical cells; fork Sc+RP distad of fork CuA_1+CuA_2 ; first crossvein r-m basad of fork MP; RP 2 branches, MP with five terminals: MP_{11} , MP_{12} , MP_2 , MP_3 , and MP_4 , fork MP_1+MP_2 basad of fork MP_3+MP_4 . Hind tibia with 2–3 lateral spines, metatibiotarsal formula: 6/7/7–8, second segment of hind tarsus with 2–3 platellae.

Male genitalia. Pygofer (Fig. 3D, E) symmetrical, dorsal margin concave and U-shaped ventrally, widened towards apex; in lateral view, lateral lobes arched extended caudally. Medioventral process triangular in ventral view. Anal segment (Fig. 3D, F) long, tubular, symmetrical, apical lobes slightly enlarged, 2.2 times longer than wide in dorsal view; anal style finger-like, not extending beyond anal segment. Gonostyli (Fig. 3D, E, G) symmetrical in ventral view; in inner lateral view, apical part extended, apical margin round. Aedeagus (Fig. 3H–K) in total with three processes. Right apex of periandrium with a medium-sized spinose process, curved and apex directed left-ventrocephalad; spinose process on left side near apical 1/3 of periandrium being the longest, gently curved from left to right side over periandrium and apex directed to the right side; base of periandrium ventrally with one small tooth. Endosoma slender, structure simple, left side near the middle with a stout and short spinose process, apex directed dorsocephalad.

Host plant. Unknown.

Distribution. China (Guizhou), Russia (Sakhalin Island).




Figure 1. Habitus **A, B** *Kuvera basarukini* Emeljanov, 1998, male **A** dorsal view **B** lateral view **C, D** *Kuvera campylostropa* sp. nov., male **C** dorsal view **D** lateral view.




Figure 2. Habitus **A, B** *Kuvera elongata* sp. nov., male **A** dorsal view **B** lateral view **C, D** *Kuvera laticeps* (Metcalf, 1936), male **C** dorsal view **D** lateral view.




Figure 3. *Kuvera basarukini* Emeljanov, 1998, male **A** head and thorax, dorsal view **B** face, ventral view **C** forewing **D** genitalia, lateral view **E** pygofer and gonostyli, ventral view **F** anal segment, dorsal view **G** gonostyli, inner lateral view **H** aedeagus, right side **I** aedeagus, left side **J** aedeagus, dorsal view **K** aedeagus, ventral view. Scale bars: 0.5 mm (**A**, **B**, **D–K**); 1.0 mm (**C**).

Remarks. This species can be distinguished from other species of the genus by the following characters: anal segment symmetrical; aedeagus with three processes: right spinose process of periandrium curved and apex directed left-ventrocephalad; left spinose process of periandrium being the longest, curved over periandrium and apex exceeding right lateral margin of periandrium; spinose process of endosoma stout and short, apex directed dorsocephalad.

Note. This species is recorded from China for the first time.

***Kuvera campylotropa* Zhi & Chen, sp. nov.**

<https://zoobank.org/6F94F366-14A5-4582-ABDD-5CBA315DDAEF>

Figs 1C, D, 4

Type material. Holotype: ♂, CHINA: Yunnan Province, Lushui City, Pianma Town (26°1'N, 98°37'E), 17 June 2011, leg. Yu-Jian Li, Jian-Kun Long; paratypes: 1♂ 1♀, same data as holotype; 6♂♂ 1♀, Guizhou Province, Daozhen County, Xiannvdong Nature Reserve (29°3'N, 107°25'E), 26 August 2004, leg. Xiang-Sheng Chen.

Description. Body length: male 5.1–6.3 mm ($N = 8$), female 6.1–6.5 mm ($N = 2$).

Coloration. General color blackish brown (Figs 1C, D, 4A, B). Eyes dark brown, ocelli yellowish brown. Vertex dark brown, pronotum dark brown and mesonotum blackish brown. Face generally blackish brown. Rostrum generally dark brown except darker tip. Forewing semi-translucent, stigma blackish brown. Hind tibiae brown and abdominal sternites blackish brown.

Head and thorax. Vertex (Fig. 4A) broad, 4.2 times wider than long; anterior margin slightly arched convex, posterior margin slightly arched concave; median carina reaching transverse carinae. Frons (Fig. 4B) 1.2 times as wide as long, median carina indistinct, extending from slightly above level of lateral ocelli to median ocellus. Clypeus with median carina distinct and elevated throughout. Pronotum (Fig. 4A) 4.4 times longer than vertex, posterior margin nearly at right angle. Mesonotum 1.9 times longer than pronotum and vertex combined. Forewing (Fig. 4C) 3.1 times longer than wide, with 11 apical and 6 subapical cells; fork Sc+RP distad of fork CuA_1+CuA_2 ; first crossvein r-m basad of fork MP; RP 3 branches, MP with five terminals: MP_{11} , MP_{12} , MP_2 , MP_3 , and MP_4 , fork MP_1+MP_2 basad of fork MP_3+MP_4 . Hind tibia with 3 lateral spines, metatibiotarsal formula: 6/7/8, second segment of hind tarsus with 3 platellae.

Male genitalia. Pygofer (Fig. 4D, E) symmetrical, dorsal margin concave and U-shaped ventrally, widened towards apex; in lateral view, lateral lobes arched extended caudally. Medioventral process triangular in ventral view. Anal segment (Fig. 4D, F) long, tubular, symmetrical, apical lobes slightly enlarged, 2.6 times longer than wide in dorsal view; anal style finger-like, not extending beyond anal segment. Gonostyli (Fig. 4D, E, G) symmetrical in ventral view; in inner lateral view, apical part extended, apical margin round. Aedeagus (Fig. 4H–K) in total with three processes. Spinose process on right side at apex of periandrium being the longest, slightly curved and apex directed left-dorsocephalad; left side in the middle with a strongly curved spinose




Figure 4. *Kuvera campylotropa* sp. nov., male **A** head and thorax, dorsal view **B** face, ventral view **C** forewing **D** genitalia, lateral view **E** pygofer and gonostyli, ventral view **F** anal segment, dorsal view **G** gonostyli, inner lateral view **H** aedeagus, right side **I** aedeagus, left side **J** aedeagus, dorsal view **K** aedeagus, ventral view. Scale bars: 0.5 mm (**A**, **B**, **D–K**); 1.0 mm (**C**).

process, apex directed ventrocephalad; base of periandrium ventrally with several small teeth. Endosoma slender, structure simple, dorsal margin near the middle with a stout and short spinose process, apex directed dorsocephalad.

Host plant. Unknown.

Distribution. China (Guizhou, Yunnan).

Etymology. The specific name is derived from the Latin *campylotropus*, meaning curved, referring to the strong curved spinose process on the left side of periandrium.

Remarks. This new species is similar to *K. ussuriensis* (Vilbaste, 1968), but differs in: (1) “right” spinose process of periandrium originating from right apex (in *K. ussuriensis*, “right” spinose process of periandrium originating from ventral apex); (2) left spinose process of periandrium shorter than right one in lateral view (the latter longer than right one); and (3) spinose process of endosoma not reaching apex of endosoma (in *K. ussuriensis*, spinose process of the endosoma nearly reaching apex of endosoma). It also closely resembles *Kuvera kurilensis* Anufriev, 1987, however, it differs in that: (1) right spinose process of periandrium longer than left one in lateral view (in *K. kurilensis*, both processes about equal in length); and (2) spinose process of endosoma not reaching apex of endosoma (in *K. kurilensis*, spinose process of endosoma nearly reaching apex of endosoma).

***Kuvera elongata* Zhi & Chen, sp. nov.**

<https://zoobank.org/0BB30DF7-8F44-4486-B262-8E069F7892E8>

Figs 2A, B, 5

Type material. *Holotype*: ♂, CHINA: Guizhou Province, Tongren City, Fanjingshan National Nature Reserve, Jinding (27°54'N, 108°42'E), 31 May 2002, leg. Xiang-Sheng Chen; *paratypes*: 8♂♂ 1♀, same data as holotype; 2♂♂, Guizhou Province, Tongren City, Fanjingshan National Nature Reserve, Yinjiang County, Yongyi Township, (27°54'N, 108°38'E), 29 May 2002, leg. Xiang-Sheng Chen.

Description. Body length: male 4.8–5.8 mm ($N = 11$), female 6.0 mm ($N = 1$).

Coloration. General color blackish brown (Figs 2A, B, 5A, B). Eyes dark brown, ocelli light yellowish. Vertex brown, pronotum dark brown and mesonotum blackish brown. Face generally blackish brown. Rostrum generally brown except darker tip. Forewing semi-translucent, stigma dark brown. Hind tibiae brown and abdominal sternites blackish brown.

Head and thorax. Vertex (Fig. 5A) broad, 2.7 times wider than long; anterior margin slightly arched convex, posterior margin arched concave; median carina reaching transverse carinae. Frons (Fig. 5B) 1.1 times as wide as long, median carina indistinct, extending from basal 1/4 to median ocellus. Clypeus with median carina distinct and elevated throughout. Pronotum (Fig. 5A) 2.3 times longer than vertex, posterior margin nearly at right angle. Mesonotum 1.6 times longer than pronotum and vertex combined. Forewing (Fig. 5C) 2.7 times longer than wide, with 11 apical and 6 sub-apical cells; fork Sc+RP distad of fork CuA_1+CuA_2 ; first crossvein r-m basad of fork




Figure 5. *Kuvera elongata* sp. nov., male **A** head and thorax, dorsal view **B** face, ventral view **C** forewing **D** genitalia, lateral view **E** pygofer and gonostyli, ventral view **F** anal segment, dorsal view **G** gonostyli, inner lateral view **H** aedeagus, right side **I** aedeagus, left side **J** aedeagus, dorsal view **K** aedeagus, ventral view. Scale bars: 0.5 mm (**A, B, D–K**); 1.0 mm (**C**).

MP; RP 3 branches, MP with five terminals: MP_{11} , MP_{12} , MP_2 , MP_3 , and MP_4 , fork MP_1+MP_2 basad of fork MP_3+MP_4 . Hind tibia with 3 lateral spines, metatibiotarsal formula: 6/7/8, second segment of hind tarsus with 2–3 platellae.

Male genitalia. Pygofer (Fig. 5D, E) symmetrical, dorsal margin concave and U-shaped ventrally, widened towards apex; in lateral view, lateral lobes arched extended caudally. Medioventral process campanulate in ventral view. Anal segment (Fig. 5D, F) long, tubular, symmetrical, apical lobes slightly enlarged, 2.0 times longer than wide in dorsal view; anal style finger-like, not extending beyond anal segment. Gonostyli (Fig. 5D, E, G) symmetrical in ventral view; in inner lateral view, apical part extended, apical margin round. Aedeagus (Fig. 5H–K) with three processes in total. Spinose process on right side at apex of periandrium being the shortest, slightly curved outward and apex directed dorsocephalad; left side near base with a slightly curved long spinose process, apex strongly recurved and directed to left side; base of periandrium ventrally with several small teeth. Endosoma slender, structure simple, left side near the middle with a stout and long spinose process, which extended beyond the apex of the endosoma, apex directed ventrad.

Host plant. Grass.

Distribution. China (Guizhou).

Etymology. The specific name is derived from the Latin *elongatus*, meaning elongated, referring to the elongated spinose process on the left side of endosoma.

Remarks. This new species is similar to *K. vilbastei* Anufriev, 1987 and *K. huoditangensis* Luo, Liu & Feng, 2019, but differs in: (1) left spinose process of periandrium not exceeding right lateral margin of periandrium (in *K. vilbastei* and *K. huoditangensis*, left spinose process of periandrium exceeding right lateral margin of periandrium); (2) spinose process of endosoma extending beyond the apex of the endosoma (spinose process of endosoma not extending beyond the apex of the endosoma in *K. vilbastei* and *K. huoditangensis*); and (3) anal segment symmetrical (asymmetrical in *K. vilbastei* and *K. huoditangensis*).

Kuvera laticeps (Metcalf, 1936)

Figs 2C, D, 6, 7

Cixius latifrons Melichar, 1902: 85, preoccupied by *Cixius latifrons* Walker, 1851.

Cixius laticeps Metcalf, 1936: 180, nom. nov. for *Cixius latifrons* Melichar, 1902.

Kuvera laticeps (Metcalf, 1936): combination by Anufriev 1987: 6.

Material examined. CHINA: 16♂♂ 22♀♀, Guizhou Province, Weining County, Xue-shan Town (27°11'N, 104°6'E), 28–29 September 2016, leg. Jian-Kun Long, Hong-Xing Li, Ya-Lin Yao.

Description. Body length: male 5.4–6.2 mm ($N=16$), female 6.1–6.8 mm ($N=22$).

Coloration. General color blackish brown (Figs 2C, D, 6A, B). Eyes brown, ocelli yellow. Vertex brown, pronotum brown and mesonotum blackish brown.




Figure 6. *Kuvera laticeps* (Metcalf, 1936), male **A** head and thorax, dorsal view **B** face, ventral view **C** forewing **D** genitalia, lateral view **E** pygofer and gonostyli, ventral view **F** anal segment, dorsal view **G** gonostyli, inner lateral view **H** aedeagus, right side **I** aedeagus, left side **J** aedeagus, dorsal view **K** aedeagus, ventral view. Scale bars: 0.5 mm (**A, B, D–K**); 1.0 mm (**C**).

Frons generally brown and clypeus blackish brown. Rostrum generally brown except darker tip. Forewing semi-translucent, with a very small irregular blackish brown spot at branch of Y-vein, stigma dark brown. Hind tibiae brown and abdominal sternites blackish brown.

Head and thorax. Vertex (Fig. 6A) broad, 3.0 times wider than long; anterior margin slightly arched convex, posterior margin arched concave; median carina reaching transverse carinae. Frons (Fig. 6B) 1.2 times as wide as long, median carina indistinct, extending from basal 1/4 to median ocellus. Clypeus with median carina distinct and elevated throughout. Pronotum (Fig. 6A) 2.8 times longer than vertex, posterior margin nearly at right angle. Mesonotum 1.6 times longer than pronotum and vertex combined. Forewing (Fig. 6C) 3.0 times longer than wide, with 11 apical and 6 sub-apical cells; fork Sc+RP distad of fork CuA₁+CuA₂; first crossvein r-m basad of fork MP; RP 3 branches, MP with five terminals: MP₁₁, MP₁₂, MP₂, MP₃, and MP₄, fork MP₁+MP₂ basad of fork MP₃+MP₄. Hind tibia with 3–4 lateral spines, metatibiotarsal formula: 6/7/8, second segment of hind tarsus with 2–4 platellae.

Male genitalia. Pygofer (Fig. 6D, E) symmetrical, dorsal margin concave and U-shaped ventrally, slightly widened towards apex; in lateral view, lateral lobes arched extended caudally. Medioventral process triangular in ventral view. Anal segment (Fig. 6D, F) long, tubular, nearly symmetrical, apical lobes slightly enlarged, 2.4 times longer than wide in dorsal view; anal style finger-like, not extending beyond anal segment. Gonostyli (Fig. 6D, E, G) symmetrical in ventral view; in inner lateral view, apical part extended, apical margin round. Aedeagus (Fig. 6H–K) in total with three processes. Spinose process on right side at apex of periandrium being the longest, gently curved from right to left side over periandrium, apex strongly recurved at 90° and directed apically; left side near base with a straight long spinose process, apex directed ventrocephalad; base of periandrium ventrally with several very small teeth. Endosoma slender, structure simple, left side near the middle with a stout and short spinose process, nearly straight, apex directed dorsocephalad.

Female genitalia. Tergite IX (Fig. 7A, B, D) moderately sclerotized, with a large wax plate, nearly oval, dorsal and ventral margins concave. Anal segment (Fig. 7C) rectangular, 2.1 times wider than long in dorsal view, anal style strap-like. Gonapophysis VIII (Fig. 7E) elongate, and slightly curved upwards. Gonapophysis IX (Fig. 7F) with two middle teeth, distance ratio between distal middle tooth to apex and length of denticulate portion is 2.6. Gonoplac (Fig. 7G) rod-like, 4.6 times longer than wide in lateral view. Posterior vagina pattern as shown in Figure 7H, I.

Host plant. Unknown.

Distribution. China (Guizhou, Sichuan).

Note. The female genitalia of this species are described and illustrated for the first time.

Remarks. This species can be distinguished from other species of the genus by the following characters: anal segment symmetrical; right spinose process of periandrium being the longest, S-shaped, curved over periandrium and apex exceeding left lateral margin of periandrium; left spinose process of periandrium straight, apex directed ventrocephalad; spinose process of endosoma stout and short straight, apex directed dorsocephalad.




Figure 7. *Kuvera laticeps* (Metcalf, 1936), female **A** genitalia, lateral view **B** genitalia, ventral view **C** anal segment, dorsal view **D** tergite IX, caudal view **E** gonapophysis VIII and gonocoxa VIII, ventral view **F** gonapophysis IX, lateral view **G** gonoplac, inner lateral view **H** posterior vagina, ventral view **I** posterior vagina, dorsal view. Scale bars: 0.5 mm.

***Kuvera ussuriensis* (Vilbaste, 1968)**

Figs 8–10

Betacixius ussuriensis Vilbaste, 1968: 9.

Kuvera ussuriensis (Vilbaste, 1968): combination by Anufriev 1987: 17.

Material examined. CHINA: 4♂♂ 7♀♀, Hebei Province, Xinglong County, Wulingshan National Nature Reserve (40°36'N, 117°29'E), 14 August 2010, leg. Li-Xia Xie, Da-Xing Yang, Rong Huang; 50♂♂ 38♀♀, Shanxi Province, Yicheng County, Lishan National Nature Reserve, Dahe Forest Farm (35°27'N, 111°56'E), 23–25 July 2012, leg. Pei Zhang; 28♂♂ 23♀♀, Shanxi Province, Qinshui County, Zhongcun Town, Zhangma Village (35°35'N, 111°57'E), 22 July 2012, leg. Pei Zhang; 26♂♂ 28♀♀, Shanxi Province, Lishan National Nature Reserve (35°23'N, 111°59'E) (1300–




Figure 8. *Kuvera ussuriensis* (Vilbaste, 1968), female **A** genitalia, lateral view **B** genitalia, ventral view **C** anal segment, dorsal view **D** tergite IX, caudal view **E** gonapophysis VIII and gonocoxa VIII, ventral view **F** gonapophysis IX, lateral view **G** gonoplac, inner lateral view **H** posterior vagina, ventral view **I** posterior vagina, dorsal view. Scale bars: 0.5 mm.

2200 m), 31 July 2012, leg. Pei Zhang; 31♂♂ 23♀♀, Shanxi Province, Lishan National Nature Reserve (1300–2200 m), 12–18 July 2012, leg. Xiao-Hui Hou; 1♂ 4♀♀, Sichuan Province, Nanchong City, Dayou Township (30°48'N, 106°41'E), 10 May 2008, leg. Zai-Hua Yang; 5♂♂ 3♀♀, Sichuan Province, Luding County, Yanzigou Town (29°42'N, 102°1'E), 11 August 2015, leg. Hong-Ping Zhan, Wen-Song Li; 2♂♂ 2♀♀, Sichuan Province, Qingchuan County, Tangjiahe National Nature Reserve (32°35'N, 104°46'E), 24 August 2007, leg. Ze-Hong Meng; 6♂♂ 10♀♀, Mianyang City, Pingwu County, Baima Tibetan Township, Wanglang Nature Reserve (32°54'N, 104°9'E); leg. Zai-Hua Yang, Wen Zhang; 2♂♂ 4♀♀, Yunnan Province, Yingjiang County, Xima Town (24°45'N, 97°42'E), 29–30 May 2011, leg. Zai-Hua Yang, Jian-Kun Long; 4♂♂, Yunnan Province, Pingbian County, Daweishan National Nature Reserve (22°56'N, 103°42'E), 20 August 2017, leg. Nian Gong; 10♂♂ 13♀♀, Yunnan Province, Xichou County, Fadou (23°23'N, 104°47'E), 28 June 2013, leg. Yan Zhi, Qiang Luo, Yong-Jin



Figure 9. Adult of *Kuvera ussuriensis* (Vilbaste, 1968), dorsal view, female (Caohai National Natural Reserve, Weining County, Guizhou Province, 2 August 2017, photograph by Xiang-Sheng Chen).

Sui; 2♂♂ 2♀♀, Yunnan Province, Maguan County, Dulong Town, Jinzhuping Village (22°56'N, 104°30'E), 13–14 August 2017, leg. Yan Zhi, Qiang Luo, Nian Gong; 8♂♂ 8♀♀, Guangxi Province, Longsheng County, Huaping National Nature Reserve (25°36'N, 109°56'E), 26 April 2012, leg. Jian-Kun Long, Zai-Hua Yang; 7♂♂ 3♀♀, Guangxi Province, Longsheng County, Huaping National Nature Reserve, 18–19 May 2012, leg. Jian-Kun Long, Zhi-Hua Fan; 9♂♂ 10♀♀, Shaanxi Province, Zhouzhi County, Houzhenzi Town (33°51'N, 107°50'E), 4–7 August 2010, leg. Pei Zhang, Zhi-Min Chang, Yan-Li Zheng, Ke-Bin Li; 5♂♂ 5♀♀, Shaanxi Province Xi'an City, Cuihuashan (33°58'N, 109°1'E), 27–28 August 2008, leg. Yu-Jian Li; 2♂♂ 2♀♀, Shaanxi Province, Taibai County (34°4'N, 107°19'E), 22 August 2016, leg. Nian Gong; 2♂♂ 5♀♀, Hunan Province, Wugang City, Yunshan National Forest Park (26°40'N, 110°37'E), May 2016, leg. Xiang-Sheng Chen; 2♂♂, Hunan Province, Yongshun County, Xiaoxi Town (28°44'N, 110°15'E), 20–21 August 2016, leg. Yong-Shun Ding, Ying-Jian Wang; 6♂♂ 2♀♀, Anhui Province, Huangshan city, Tangkou town (30°4'N, 118°11'E) (500m), 20 May 2008, leg. Zheng-Guang Zhang; 22♂♂ 33♀♀, Guizhou Province, Weining County, Caohai National Nature Reserve (26°52'N, 104°14'E) (2171 m), 1–5 August 2017, leg. Caohai Collection Team; 1♂ 5♀♀, Guizhou Province, Weining County, Xueshan Town, Zhuopu Village (27°11'N, 104°6'E), 21 August 1986, leg. Zi-Zhong Li; 6♂♂ 17♀♀, Guizhou Province, Daozhen County, Xiannvdong (29°3'N, 107°25'E), 29–31 May 2004, leg. Bin Zhang, Pian Xu; 25♂♂ 44♀♀, Guizhou Province, Daozhen



Figure 10. The habitat of *Kuvera ussuriensis* (Vilbaste, 1968) (3 August 2017, Caohai National Natural Reserve, Weining County, Guizhou Province, photograph by Yan Zhi).

County, Sanqiao Town (29°3'N, 107°30'E), 22–24 May 2004, leg. Xiang-Sheng Chen, Bin Zhang, Pian Xu; 4♂♂ 17♀♀, Guizhou Province, Daozhen County, Dashahe Nature Reserve (29°9'N, 107°36'E), 29–31 May 2004, leg. Xiang-Sheng Chen; 2♂♂ 3♀♀, Guizhou Province, Daozhen County, Dashahe Nature Reserve, 20 August 2004, leg. Xiang-Sheng Chen; 3♂♂, Guizhou Province, Luodian County, Luosha Township, Zheren Village (25°41'N, 106°36'E), 9 May 2013, leg. Jian-Kun Long; 6♂♂ 14♀♀, Guizhou Province, Anlong County (25°5'N, 105°29'E), 27 August 2012, leg. Jian-Kun Long, Wei-Bin Zheng, Shi-Yan Xu; 8♂♂ 8♀♀, Guizhou Province, Suiyang County, Kuankuoshui National Nature Reserve (28°14'N, 107°12'E), 2–4 June 2010, leg. Yan-Li Zheng; 10♂♂ 6♀♀, Guizhou Province, Tongren City, Fanjingshan National Nature Reserve (27°55'N, 108°42'E), 20–24 September 2011, leg. Wei-Bin Zheng, Zhi-Min Chang, Xiao-Fei Yu, Zhi-Hua Fan; 17♂♂ 4♀♀, Guizhou Province, Tongren City, Fanjingshan National Nature Reserve, Yinjiang County, Yongyi Township, Dayuanzhi Village, (27°54'N, 108°38'E), 29 May 2002, leg. Xiang-Sheng Chen; 1♂, Guizhou Province, Tongren City, Fanjingshan National Nature Reserve, Heihewan (27°50'N, 108°46'E), 30 July 2014, leg. Meng-Shu Dong; 1♀, Guizhou Province, Tongren City, Fanjingshan National Nature Reserve, Jinding (27°54'N, 108°42'E), 1 September 1996, leg. Mao-Fa Yang; 3♀♀, Guizhou Province, Tongren City, Fanjingshan National Nature Reserve, Jinding, 30 July 2001, leg. Mao-Fa Yang, Guo-Dong Ren; 4♂♂ 6♀♀, Guizhou Province, Leishan County, Leigongshan National Forest Park (26°21'N,




Figure 11. Distribution records of species from the genus *Kueneria*: *K. amurensis* (red triangle); *K. basarukini* (pink triangle); *K. brunnetii* (green hexagon); *K. brunnea* (red square); *K. campylotropa* sp. nov. (blue hexagon); *K. communis* (red circle); *K. elongata* sp. nov. (yellow inverse triangle); *K. flaviceps* (green inverse triangle); *K. hagibasanensis* (red star); *K. hallasanensis* (yellow triangle); *K. hama* (green triangle); *K. huoditangensis* (red inverse triangle); *K. kurilensis* (red hexagon); *K. latticeps* (pink hexagon); *K. ligustri* (yellow hexagon); *K. longipennis* (yellow star); *K. longwangshanensis* (yellow square); *K. pallidula* (blue triangle); *K. semihyalina* (blue square); *K. similis* (blue circle); *K. taiwana* (pink inverse triangle); *K. tappanella* (yellow circle); *K. toroensis* (green star); *K. transversa* (blue star); *K. ussuriensis* (green circle); *K. vilbastei* (blue inverse triangle); and *K. yecheonensis* (pink circle).

108°9'E), 4–6 July 2011, leg. Wei-Bin Zheng, Jian-Kun Long; 3♂♂ 2♀♀, Guizhou Province, Leishan County, Leigongshan National Forest Park, Lianhuaping, 31 May–3 June 2005, leg. Zi-Zhong Li, Qiong-Zhang Song, Bin Zhang; 5♂♂ 4♀♀, Guizhou Province, Duyun City, Chachang (26°24'N, 107°36'E), 12 May 2014, leg. Ming Ning, Gai-Ping Yang, Ding-Guo Li; 2♂♂ 1♀, Guizhou Province, Duyun City, Chachang, 16 August 2014, leg. Gai-Ping Yang, Ding-Guo Li; 1♂ 4♀♀, Guizhou Province, Wangmo County, Dayi Town (25°10'N, 106°06'E), 22 August 2012, leg. Shi-Yan Xu, Wei-Bin Zheng; 1♂ 1♀, Guizhou Province, Guiyang City, Guizhou Botanical Garden (26°37'N, 106°44'E), 18 June 2008, leg. Jun-Qiang Ni; 3♂♂ 3♀♀, Guizhou Province, Guiyang City, Wudang District (26°38'N, 106°45'E), 5 June 2009, leg. Qiong-Zhang Song; 1♂ 2♀♀, Guizhou Province, Zunyi City, Loushanguan (28°1'N, 106°51'E), 21 September 2017, leg. Bin Yan; 15♂♂ 3♀♀, Guizhou Province, Xishui County, Linjiang (28°19'N, 106°12'E), 1 June 2006, leg. Xiang-Sheng Chen.

Supplementary description. Female genitalia. Tergite IX (Fig. 8A, B, D) moderately sclerotized, with a large wax plate, nearly oval, dorsal and ventral margins concave. Anal segment (Fig. 8C) rectangular, 1.8 times wider than long in dorsal view, anal style strap-like. Gonapophysis VIII (Fig. 8E) elongate, and slightly curved upwards. Gonapophysis IX (Fig. 8F) with two middle teeth, distance ratio between distal middle tooth to apex and length of denticulate portion is 2.9. Gonoplac (Fig. 8G) rod-like, 4.0 times longer than wide in lateral view. Posterior vagina pattern as shown in Figure 8H, I.

Host plant. *Artemisia mongolica* (Fisch. ex Bess.) Nakai (Asteraceae) (Fig. 10).

Distribution. China (Anhui, Guangxi, Guizhou, Hebei, Hunan, Shaanxi, Shanxi, Sichuan, Yunnan), Japan, Russia.

Remarks. This species can be distinguished from other species of the genus by the following characters: anal segment symmetrical; aedeagus with three processes: left spinose process of periandrium long, gently curved and apex directed left-ventrocephalad; ventral surface of periandrium with a spinose process, slightly curved and apex directed right-cephalad; spinose process of endosoma stout and long, nearly reaching apex of endosoma.

Note. The female genitalia of this species are described and illustrated for the first time.

Discussion

The biology of few *Kuvera* species throughout the world are well-known. The plant associations of genus have been reported in several previous studies (Anufriev 1987; Emeljanov 2015; Luo et al. 2019). In this study, we found *Kuvera ussuriensis* (Vilbaste, 1968) on *Artemisia mongolica* (Fisch. ex Bess.) Nakai.

Based on data from published information and our field surveys, the distribution records of all twenty-seven known species of *Kuvera* was summarized in Figure 11. Up to now, the genus presents a distribution pattern in the Palearctic, Sino-Japanese, and Oriental regions. Compared with Luo et al. (2019), new distribution records of several species have been added recently (Luo et al. 2022 and this study). We believe that the

actual distribution range of most species is still unclear. *Kuvera brunettii* Muir, 1922, *K. brunnea* (Dlabola, 1957), *K. hagilsanensis* Rahman, Kwon & Suh, 2017, *K. hal-lasanensis* Rahman, Kwon & Suh, 2017, *K. ligustri* Matsumura, 1914, *K. longipennis* Matsumura, 1914 and *K. longwangshanensis* Luo, Liu & Feng, 2019 are known only from the type locality, and further collecting and investigation of this genus are still needed. The complex and variable geomorphological environment and rich biological resources of the distribution area create a variety of habitat types, which are likely reasons for the rich species diversity of the genus *Kuvera*.

Acknowledgements

The authors are grateful to the specimen collectors for their hard work in the field collections. We wish to express our sincere thanks to Prof. A. Emeljanov (Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia) for providing related literature and Dr Wei Du (College of Life Sciences, Wuhan University) for identifying the host plant. This work was supported by the National Natural Science Foundation of China (No. 32060343, 31472033), the Science and Technology Support Program of Guizhou Province (grant no. 20201Y129), the Program of Excellent Innovation Talents, Guizhou Province (No. 20154021), the New Academic Talent Program of Guizhou Medical University (No. 19NSP066) and Doctoral Scientific Research Foundation of Guizhou Medical University (No. J[2020]019).

References

- Anufriev GA (1987) Review of the cixiid genus *Kuvera* Distant (Homoptera, Auchenorrhyncha, Cixiidae). In: Kapustina OG (1987) [Taxonomy of the insects of Siberia and Soviet Far East], *Taksonomiia nasekomykh Sibiri i Dal'nego Vostoka SSR.*, Vladivostok Dalnauka, Russia, 4–21.
- Anufriev GA, Emeljanov AF (1988) Suborder Cicadinea (Auchenorrhyncha). In: Lehr PA (Ed.) *Keys to the Insects of the Far East of the USSR (Vol. 2). Homoptera and Heteroptera.* Nauka Publishing House, Leningrad, 496 pp.
- Bourgoin T (1987) A new interpretation of the homologies of the Hemiptera male genitalia, illustrated by the Tettigometridae (Hemiptera, Fulgoromorpha). *Proceedings 6th Auchenorrhyncha Meeting, Turin, Italy, 7–11 September 1987*, 113–120.
- Bourgoin T (1993) Female genitalia in Hemiptera Fulgoromorpha, morphological and phylogenetic data. *Annales de la Société Entomologique de France* 29(3): 225–244.
- Bourgoin T (2022) FLOW (Fulgoromorpha Lists On the Web): A knowledge and a taxonomy database dedicated to planthoppers (Insecta, Hemiptera, Fulgoromorpha, Fulgoroidea). Version 8, updated 03 February 2022. <https://flow.hemiptera-databases.org/flow/> [Accessed on: 16 March 2022]
- Bourgoin T, Wang RR, Asche M, Hoch H, Soulier-Perkins A, Stroiński A, Yap S, Szvedo J (2015) From micropterism to hyperpterism: recognition strategy and standardized homology-driv-

- en terminology of the forewing venation patterns in planthoppers (Hemiptera: Fulgoromorpha). *Zoomorphology* 134(1): 63–77. <https://doi.org/10.1007/s00435-014-0243-6>
- Distant WL (1906) Rhynchota. III. (Heteroptera Homoptera). In: Bingham CT (Ed.) The fauna of British India including Ceylon and Burma. Taylor & Francis, London, 503 pp. <https://doi.org/10.1080/00222930608562592>
- Dlabola J (1957) Die Zikaden Afghanistans (Homopt.-Auchenorrhyncha) nach den Ergebnissen der von Herrn J. Klapperich in den Jahren 1952–1953 nach Afghanistan unternommenen Expedition. *Mitteilungen der Münchner Entomologischen Gesellschaft*, E V. München 47(5): 265–303.
- Emeljanov AF (1998) Contribution to the knowledge of the genus *Kuvera* Distant (Homoptera: Cixiidae). *Zoosystematica Rossica* 7(1): 133–137.
- Emeljanov AF (2002) Contribution to classification and phylogeny of the family Cixiidae (Hemiptera, Fulgoromorpha). *Denisia* 04: 103–112.
- Emeljanov AF (2015) Planthoppers of the family Cixiidae of Russia and adjacent territories. *Key to the fauna of Russia* 177: 1–252.
- Holt BG, Lessard JP, Borregaard MK, Fritz SA, Araújo MB, Dimitrov D, Fabre PH, Graham CH, Graves GR, Jönsson KA, Nogués-Bravo D, Wang Z, Whittaker RJ, Fjeldså J, Rahbek C (2013) An update of Wallace's zoogeographic regions of the world. *Science* 339(6115): 74–78. <https://doi.org/10.1126/science.1228282>
- Holzinger WE, Emeljanov AF, Kammerlander I (2002) The family Cixiidae Spinola, 1839 (Hemiptera: Fulgoromorpha)-a Review. *Denisia* 4: 113–138.
- Luo Y, Liu J, Feng J (2019) Two new species in the genus *Kuvera* Distant, 1906 (Hemiptera, Cixiidae, Cixiinae) from China. *ZooKeys* 832: 135–152. <https://doi.org/10.3897/zookeys.832.30301>
- Luo Y, Bourgoïn T, Zhang JL, Feng JN (2022) Distribution patterns of Chinese Cixiidae (Hemiptera, Fulgoroidea), highlight their high endemic diversity. *Biodiversity Data Journal* 10: e75303. <https://doi.org/10.3897/BDJ.10.e75303>
- Matsumura S (1914) Die Cixiinen Japans. *Annotationes Zoologicae Japonenses* 8: 393–434.
- Melichar L (1902) Homopteren aus West-China, Persien und dem Süd-Ussuri-Gebiete. *Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St.-Petersbourg*. Saint Petersburg 7: 76–146.
- Metcalf ZP (1936) Part 2. Cixiidae. In: Metcalf ZP (Ed.) *General Catalogue of the Homoptera*. Fascicule IV, North Carolina State College, Raleigh, 267 pp.
- Muir FAG (1922) New Indian Homoptera. *Records of the Indian Museum* 24: 343–355. <https://doi.org/10.26515/rzsi/v24/i3/1922/162723>
- Rahman MA, Kwon YJ, Suh SJ (2017) Three new species of the genus *Kuvera* Distant (Hemiptera: Fulgoromorpha: Cixiidae) from Korea. *Oriental Insects* 52(1): 66–78. <https://doi.org/10.1080/00305316.2017.1344741>
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <https://www.simplemappr.net> [Accessed on: 12 March 2022]
- Tsaur SC, Hsu TC, Van Stalle J (1991) Cixiidae of Taiwan, Part V. Cixiini except Cixius. *Journal of Taiwan Museum* 44(1): 1–78.
- Vilbaste J (1968) On the Cicadine Fauna of the Primorsk Region. Valgus Publishing, Tallinn, 195 pp. [In Russian]