

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/256613212>

Trigonocranus emmeae Fieber, 1876 (Hemiptera, Fulgoromorpha, Cixiidae) – a new species for Poland

Article in *ZooKeys* · July 2013

DOI: 10.3897/zookeys.319.4278 · Source: PubMed

CITATIONS

5

READS

161

5 authors, including:



Krzysztof Musik

University of Silesia in Katowice

7 PUBLICATIONS 37 CITATIONS

[SEE PROFILE](#)



Marcin Walczak

University of Silesia in Katowice

79 PUBLICATIONS 190 CITATIONS

[SEE PROFILE](#)



Łukasz Depa

University of Silesia in Katowice

137 PUBLICATIONS 353 CITATIONS

[SEE PROFILE](#)



Łukasz Junkiert

University of Silesia in Katowice

32 PUBLICATIONS 174 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Use of Scanning Electron Microscopy (SEM) in Zoology – observation of claws and pad structures of insects of family Heteropterygidae. [View project](#)



The description of nymphs of stick insects of family Heteropterygidae. [View project](#)

***Trigonocranus emmeae* Fieber, 1876 (Hemiptera, Fulgoromorpha, Cixiidae) – a new species for Poland**

Krzysztof Musik¹, Marcin Walczak¹, Łukasz Depa¹,
Łukasz Junkiert¹, Anna Jedynowicz²

¹ Department of Zoology, Faculty of Biology and Environmental Protection, University of Silesia, Bankowa 9, 40-007 Katowice, Poland ² Brzezinka, ul. Skotnicka 3A 32-600 Oświęcim, Poland

Corresponding author: Krzysztof Musik (krzysztof.musik@us.edu.pl)

Academic editor: S. Grozeva | Received 14 November 2012 | Accepted 22 February 2013 | Published 30 July 2013

Citation: Musik K, Walczak M, Depa Ł, Junkiert Ł, Jedynowicz A (2013) *Trigonocranus emmeae* Fieber, 1876 (Hemiptera, Fulgoromorpha, Cixiidae) – a new species for Poland. In: Popov A, Grozeva S, Simov N, Tasheva E (Eds) Advances in Hemipterology. ZooKeys 319: 249–253. doi: 10.3897/zookeys.319.4278

Abstract

A single macropterous female of *Trigonocranus emmeae* Fieber, 1876 has been found during the faunistic studies in semi-natural plant communities of Oświęcim city in southern Poland. It is the first record of this species in Poland. *Trigonocranus emmeae* is rarely collected within the wide range of its distribution, mostly due to its hidden life mode.

Keywords

Faunistics, new records, *Trigonocranus*, Cixiidae, Fulgoromorpha

Introduction

The Fulgoromorpha and Cicadomorpha are represented in Europe by 2053 species (Hoch 2012). In Poland the fauna of Auchenorrhyncha comprises 537 recorded species, including the latest checklist by Chudzicka (2004) and species that were missed (Łabanowski and Soika 1997, Gębicki 2003, Świerczewski and Gebicki 2003) or published later than 2004 (Gaj et al. 2009, Świerczewski and Stroński 2011a, 2011b, Świerczewski and Walczak 2011, Walczak et al. 2012).

During the recent studies in Brzezinka, a suburban district of Oświęcim city (southern Poland) a single female of *Trigonocranus emmeae* Fieber, 1876 has been collected. It is the first record of this poorly studied representative of Cixiidae in Poland.

Material and methods

Faunistic studies on planthoppers and leafhoppers (Cicadomorpha and Fulgoromorpha) were conducted during the vegetation season of 2008. The area of the research was a suburban district of Oświęcim city – Brzezinka (UTM:CA64) (50°2'51"N, 19°9'38"E).

Insects were collected by a standard sweeping net (Ø 35cm) from the end of April till the end of October, altogether 15 samples per plot were taken. The collected material was transferred to a container with ethyl acetate. In the laboratory the collected insects were mounted on glue boards and determined. The key used to identify the species was Biedermann and Niedringhaus (2004).

Chorological and ecological data used in this work are accordant to Nickel and Remane (2002) and Nickel (2003). To determine the plant associations the key provided by Matuszkiewicz (2008) was applied.

Collected material is deposited in the Collection of Department of Zoology, University of Silesia, Katowice.

Results

Trigonocranus emmeae Fieber, 1876 – new species for Polish fauna (Fig. 1).

Material examined

A single specimen of this species was collected on 22.06.2008 in Oświęcim-Brzezinka (southern Poland), [UTM CA64], 50°2'51"N, 19°9'38"E, in the meadow belonging to the *Molinio-Arrhenatheretea* class, macropterous ♀, leg. A. Jedynowicz, det. M. Walczak, C. Gębicki rev. (specimen deposited in the collection of Department of Zoology, University of Silesia).

The first locality of *T. emmeae* in Poland was located in Oświęcim-Brzezinka, Leśna street (Fig. 2). It was a *Molinio-Arrhenatheretea* class meadow surrounded by forest. The dominant plant species were: *Holcus mollis*, *Dactylis glomerata*, *Alopecurus pratensis* and *Agrostis capillaris*, in less percentage: *Anthoxanthum odoratum*, *Elymus repens*, *Carex hirta*, *C. acutiformis* and *C. vulpia*. During the studies in 2008 apart from *T. emmeae* there were 36 other species recorded in this plot. The dominant species were: *Cicadella viridis* (Linnaeus, 1758) (23.57% of collected material), *Cicadula quadrinotata* (Fabricius, 1794) (8.02%), *Arthaldeus pascuellus* (Fallén, 1826) (5.02%) and *Stenocranus major* (Kirschbaum, 1868) (3.31%) (Jedynowicz, 2009).



Figure 1. *Trigonocranus emmeae* macropterous female (photo by A. Stroński).



Figure 2. Habitat of *T. emmeae*, Oświęcim-Brzezinka (photo by A. Jedynowicz).

Discussion

According to the literature *T. emmeae* is widely distributed in Europe, but very rarely collected. This species was recorded in Austria, Switzerland, France, Great Britain (Nast 1972), Slovenia (Holzinger and Seljak 2001, Seljak et al. 2003), Croatia, Spain, Italy, Bulgaria, Germany, Georgia, southern Russia (Holzinger et al. 2003, Nickel 2003), Czech Republic (Malenovský and Lauterer 2010, 2012), Luxemburg (Niedringhaus et al. 2010) and Sweden (Larsson 2010).

According to Nickel (2003) nymphs and brachypterous adults are unpigmented with reduced number of ommatidia and live on the soil surface and leaf litter. This species oc-

curs from the end of May till the end of July, hibernates in the egg stage and is univoltine. *T. emmeae* represents the European chorological element. It lives on moderately warm sites covered by vegetation of medium density, probably feeding on roots of shrubs. The mentioned specimen was collected on a moderately moist meadow. The data from Great Britain indicates that this species may be also collected in damp sites (Bantock 2012). The host plant species is unknown (Nickel and Remane 2002, Nickel 2003). There are no data about the trophic relations of this species, therefore it is unknown if it is mono-, oligo- or most probably polyphagous. The majority of collected specimens were caught by sweeping net or Malaise traps during the dispersal flight (Nickel 2003). *T. emmeae* can be also and effectively collected by an underground pitfall trap (M. Wilson, pers. comm.).

The difficulties in collecting *T. emmeae* result in lack of detailed biological and ecological data. This species is probably not as rare as it seems, however the right collection method must be applied to reveal its cryptic presence. Distribution of this species in Poland needs further research.

Acknowledgments

We are grateful to Adam Stroiński for the photograph of the specimen. We are also grateful to the anonymous reviewers for their efforts in improving the manuscript.

References

- Bantock T (2012) An unusual record of *Trigonocranus emmeae*. Leafhopper Distribution Recording & Analysis, Newsletter of the Auchenorrhyncha recording Scheme for Britain & Ireland: 8 p.
- Biedermann R, Niedringhaus R (2004) Die Zikaden Deutschlands. Bestimmungstabellen für alle Arten. Wissenschaftlich Akademischer Buchvertrieb-Fründ, 409 pp.
- Chudzicka E (2004) Piewiki (Auchenorrhyncha = Cicadomorpha + Fulgoromorpha). In: Bogdanowicz W, Chudzicka E, Pilipiuk I, Skibińska E (Eds) Fauna Polski – charakterystyka i wykaz gatunków. Tom I. Muzeum i Instytut Zoologii PAN, Warszawa: 178–192.
- Gaj D, Walczak M, Wojciechowski W (2009) Communities of planthoppers and leafhoppers (Hemiptera: Fulgoromorpha et Cicadomorpha) inhabiting selected plant communities in the Stołowe Mountains National Park. Annals of the Upper Silesian Museum in Bytom Natural History, The Monograph 19: 1–174.
- Gębicki C (2003) Nowe stanowisko *Utecha lugens* (Germar, 1821) w Polsce (Hemiptera, Ulopidae). Acta entomologica silesiana, Vol. 9–10: 23–26.
- Hoch H (2012) Fauna Europaea: Fulgoromorpha, Cicadomorpha. Fauna Europaea version 2.5, <http://www.faunaeur.org>
- Holzinger WE, Kammerlander I, Nickel H (2003) The Auchenorrhyncha of Central Europe. Die Zikaden Mitteleuropas. Vol. 1. Fulgoromorpha, Cicadomorpha (excl. Cicadellidae). Brill Academic Publishers, Leiden – Boston, 673 pp.

- Holzinger W, Seljak G (2001) New records of planthoppers and leafhoppers from Slovenia, with a checklist of hitherto recorded species (Hemiptera, Auchenorrhyncha). *Acta Entomologica Slovenica* 9(1): 39–66.
- Jedynowicz A (2009) Piewiki (Fulgoromorpha et Cicadomorpha) okolic Oświęcimia. Master degree work, manuscript: 97 pp.
- Larsson S (2010) Spår till mångfald. Banverket, Expert och utveckling, Borlänge: 37 p.
- Łabanowski G, Soika G (1997) Nowe i mniej znane szkodniki występujące na drzewach i krzewach ozdobnych. *Progress in Plant Protection / Postępy w Ochronie Roślin*, Vol. 31(1): 218–223.
- Malenovský I, Lauterer P (2010) Additions to the fauna planthoppers and leafhoppers (Hemiptera: Auchenorrhyncha) of the Czech Republic. *Acta Musei Moraviae, Scientiae biologicae*, Brno, 95(1): 41–122.
- Malenovský I, Lauterer P (2012) Leafhoppers and planthoppers (Hemiptera: Auchenorrhyncha) of the Bílé Karpaty Protected Landscape Area and Biosphere Reserve (Czech Republic). *Acta Musei Moraviae, Scientiae biologicae*, Brno 96(2): 155–322.
- Matuszkiewicz W (2008) Przewodnik do oznaczania zbiorowisk roślinnych Polski. Wydawnictwo Naukowe PWN. Warszawa, 537 pp.
- Nast J (1972) Palearctic Auchenorrhyncha (Homoptera). An annotated check list. PWN, Warszawa: 550 pp.
- Nickel H, Remane R (2002) Artenliste der Zikaden Deutschlands, mit Angabe von Nährpflanzen, Nahrungsbreite, Lebenszyklus, Areal und Gefährdung (Hemiptera, Fulgoromorpha et Cicadomorpha). *Beiträge zur Zikadenkunde* 5: 27–64.
- Nickel H (2003) The Leafhoppers and Planthoppers of Germany (Hemiptera, Auchenorrhyncha): Patterns and strategies in a highly diverse group of phytophagous insects. Series Faunistica 28. Pensoft Publishers, Sofia-Moscow, Goecke & Evers, Keltern, 460 pp.
- Niedringhaus R, Biedermann R, Nickel H (2010) Verbreitungsatlas der Zikaden des Großherzogtums Luxemburg – Atlasband. Ferrantia, Musée national d'histoire naturelle, Luxembourg, 61: 1–395.
- Seljak G, Matis G, Miklavc J, Beber K (2003) Identifikacija potencialnih naravnih prenašalcev trsnih rumenic v podravski vinorodni deželi. Zbornik predavanj in referatov 6.slovenckega postvetovanja o varstvu rastlin, Zreče 4–6 marec 2003, 283–288.
- Świerczewski D, Gębicki C (2003) Nowe i rzadkie gatunki piewików w faunie Polski (Hemiptera: Fulgoromorpha et Cicadomorpha). *Acta entomologia silesiana*, Bytom 11 (1–2), 63–73.
- Świerczewski D, Stroiński A (2011a) The first record of the Nearctic treehopper *Stictocephala bisonia* in Poland (Hemiptera: Cicadomorpha: Membracidae) with some comments on this potential pest. *Polish Journal of Entomology* 80: 13–22. doi: 10.2478/v10200-011-0002-7
- Świerczewski D, Stroiński A (2011b) Planthoppers and leafhoppers of the Przedborski Landscape Park (Hemiptera: Fulgoromorpha et Cicadomorpha). *Polish Journal of Entomology*, 80: 277–290. doi: 10.2478/v10200-011-0020-5
- Świerczewski D, Walczak M (2011) New records of leafhoppers for Poland (Hemiptera: Cicadomorpha). *Polish Journal of Entomology* 80: 291–298. doi: 10.2478/v10200-011-0021-4
- Walczak M, Musik K, Mokrzycka A (2012) *Japananus hyalinus* (Osborn, 1900) – a new leafhopper for Polish fauna (Hemiptera: Cicadomorpha). *Wiadomości Entomologiczne* 31 (4): 242–250.